

AMENDMENTS TO THE CLAIMS

- 1.-2. (canceled)
3. (currently amended) A pharmaceutical composition ~~as recited in claim 2~~ comprising:  
a. a therapeutically effective amount of a first compound, said first compound being an estrogen agonist/antagonist; and  
b. a therapeutically effective amount of a second compound, said second compound being a prostaglandin or a prostaglandin agonist/antagonist and a pharmaceutical carrier;  
wherein the estrogen agonist/antagonist is droloxifene, raloxifene, tamoxifen, 4-hydroxy-tamoxifen,  
*Cis*-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;  
(-)-*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;  
*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;  
*Cis*-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydro-naphthalene;  
1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;  
*Cis*-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or  
1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.
4. (original) A pharmaceutical composition according to claim 3 wherein the second compound is PGD<sub>1</sub>, PGD<sub>2</sub>, PGE<sub>2</sub>, PGE<sub>1</sub>, PGF<sub>2</sub>, PGF<sub>2</sub>α or 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(1H-tetrazol-5-yl)-hexyl]-cyclopentanone.
5. (canceled)
6. (previously presented) A pharmaceutical composition according to claim 4 wherein the second compound is PGE<sub>2</sub>.
7. (previously presented) A pharmaceutical composition according to claim 4 wherein the second compound is 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(2H-tetrazol-5-yl)-hexyl]-cyclopentanone.

8. (original) A pharmaceutical composition according to claim 4 wherein the estrogen agonist/antagonist is
- Cis*-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- (-)-*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis*-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydrohaphthalene;
- 1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;
- Cis*-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or
- 1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.
9. (original) A pharmaceutical composition according to claim 8 wherein the second compound is PGE<sub>2</sub>.
10. (original) A pharmaceutical composition according to claim 8 wherein the second compound is 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(2H-tetrazol-5-yl)-hexyl]-cyclopentanone.
11. (currently amended) A method for treating a mammal having a condition which presents with low bone mass comprising administering to a mammal having a condition which presents with low bone mass
- a. a therapeutically effective amount of a first compound, said first compound being an estrogen agonist/antagonist; and
- b. a therapeutically effective amount of a second compound, said second compound being a prostaglandin or a prostaglandin agonist/antagonist;
- wherein the estrogen agonist/antagonist is droloxifene, raloxifene, tamoxifen, 4-hydroxy-tamoxifen, idoxifene, centrachroman,
- Cis*-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- (-)-*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

Cis-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydro-naphthalene;

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;

Cis-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or

1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline and wherein the second compound is PGD<sub>1</sub>, PGD<sub>2</sub>, PGE<sub>2</sub>, PGE<sub>1</sub>, PGF<sub>2</sub>, PGF<sub>2</sub>α or 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(1H-tetrazol-5-yl)-hexyl]-cyclopentanone.

12. (currently amended) A method as recited in claim 11 wherein the estrogen agonist/antagonist is droloxifene, raloxifene, tamoxifen, 4-hydroxy-tamoxifen, idoxifene, centrachroman,

Cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

(-)-Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

Cis-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydro-naphthalene;

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;

Cis-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or

1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.

13. (currently amended) A method as recited in claim 11 wherein the second compound is PGD<sub>1</sub>, PGD<sub>2</sub>, PGE<sub>2</sub>, PGE<sub>1</sub>, PGF<sub>2</sub>, PGF<sub>2</sub>α or 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(1H-tetrazol-5-yl)-hexyl]-cyclopentanone.
14. (original) A method as recited in claim 13 wherein the estrogen agonist/antagonist is droloxifene.
15. (canceled)
16. (original) A method as recited in claim 14 wherein the second compound is 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(1H-tetrazol-5-yl)-hexyl]-cyclopentanone.

17. (original) A method as recited in claim 14 wherein the condition which presents with low bone mass is osteoporosis.
18. (original) A method as recited in claim 13 wherein the estrogen agonist/antagonist is  
*Cis*-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;  
(-)-*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;  
*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;  
*Cis*-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydro-naphthalene;  
1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;  
*Cis*-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or  
1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.
19. (original) A method as recited in claim 18 wherein the second compound is PGE<sub>2</sub>.
20. (original) A method as recited in claim 18 wherein the second compound is 3S-(3-Hydroxy-4-phenyl-butyl)-2R-[6-(1H-tetrazol-5-yl)-hexyl]-cyclopentanone.
21. (original) A method as recited in claim 18 wherein the condition which presents with low bone mass is osteoporosis.
22. (original) A method as recited in claim 14 wherein the first compound and the second compounds are administered substantially simultaneously.
23. (original) A method as recited in claim 14 wherein the second compound is administered for a period of from about three months to about three years.
24. (original) A method as recited in claim 23 followed by administration of the first compound for a period of from about three months to about three years without the administration of the second compound during the period of from about three months to about three years.
25. (original) A method as recited in claim 23 followed by administration of the first compound for a period greater than about three years without the administration of the second compound during the greater than about three year period.
26. (original) A method as recited in claim 18 wherein the first compound and the second compounds are administered substantially simultaneously.

27. (original) A method as recited in claim 18 wherein the second compound is administered for a period of from about three months to about three years.
28. (original) A method as recited in claim 27 followed by administration of the first compound for a period of from about three months to about three years without the administration of the second compound during the period of from about three months to about three years.
29. (original) A method as recited in claim 27 followed by administration of the first compound for a period greater than about three years without the administration of the second compound during the greater than about three year period.
- 30-45. (canceled)
46. (previously presented) A pharmaceutical composition comprising:
- a. a therapeutically effective amount of a first compound, said first compound being
- Cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- (-)-Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydro-naphthalene;
- 1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;
- Cis-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or
- 1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline; and
- b. a therapeutically effective amount of a second compound, said second compound being sodium fluoride, a parathyroid hormone, or growth hormone.
47. (original) A pharmaceutical composition as recited in claim 46 additionally comprising a pharmaceutical carrier.
48. (original) A pharmaceutical composition as recited in claim 47 wherein the second compound is sodium fluoride.

49. (original) A pharmaceutical a composition as recited in claim 47 wherein the second compound is a parathyroid hormone.
50. (original) A pharmaceutical composition as recited in claim 47 wherein the second compound is growth hormone.
51. (canceled)
52. (previously presented) A method for treating a mammal having a condition which presents with low bone mass comprising administering to a mammal having a condition which presents with low bone mass
- a. a therapeutically effective amount of a first compound, said first compound being
- Cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- (-)-Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
- Cis-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydrohaphthalene;
- 1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;
- Cis-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or
- 1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline; and
- b. a therapeutically effective amount of a second compound, said second compound being sodium fluoride, a parathyroid hormone, or growth hormone.
53. (original) A method as recited in claim 52 wherein the second compound is sodium fluoride.
54. (original) A method as recited in claim 52 wherein the second compound is a parathyroid hormone.
55. (original) A method as recited in claim 52 wherein the second compound is growth hormone.
56. (canceled)
57. (original) A method as recited in claim 52 wherein the condition which presents with low bone mass is osteoporosis.

58. (original) A method as recited in claim 52 wherein the first compound and the second compound are administered substantially simultaneously.
59. (original) A method as recited in claim 52 wherein the second compound is administered for a period of from about three months to about three years.
60. (original) A method as recited in claim 59 followed by administration of the first compound for a period of from about three months to about three years without the administration of the second compound during the period of from about three months to about three years.
61. (original) A method as recited in claim 59 followed by administration of the first compound for a period greater than about three years without the administration of the second compound during the greater than about three year period.
62. (original) A method for treating mammals which present with low bone mass comprising administering to a mammal having a condition which presents with low bone mass the pharmaceutical composition of claim 46.
- 63-64. (canceled)
65. (previously presented) A kit containing a treatment for a condition which presents with low bone mass comprising:
- a. a therapeutically effective amount of
    - Cis*-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
    - (-)-*Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
    - Cis*-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;
    - Cis*-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydro-naphthalene;
    - 1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;
    - Cis*-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol; or
    - 1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline and a pharmaceutically acceptable carrier in a first unit dosage form;
  - b. a therapeutically effective amount of sodium fluoride, a parathyroid hormone, or growth hormone and a pharmaceutically acceptable carrier in a second unit dosage form; and

- c. container means for containing said first and second dosage forms.
66. (original) The pharmaceutical composition as recited in claim 34 wherein the first compound is droloxifene.
67. (original) The method as recited in claim 36 wherein the first compound is droloxifene.
68. (original) The method as recited in claim 40 wherein the first compound is droloxifene.
69. (original) The method as recited in claim 41 wherein the first compound is droloxifene.
- 70-71. (canceled)
72. (original) The kit as recited in claim 45 wherein the first compound is droloxifene.
- 73-108. (canceled)